

Wild Rice Study - Hydroponic Tests
Summary of dose response analysis to date
Emily Peters
3/10/2014 – corrected plots against mean initial sulfide

Analysis description

Data from 3 hydroponic tests (Range Finder, Definitive 1, and Definitive 2) were independently fit to a 4-parameter logistic model using R software. These analyses use growth change as the response variable, which provides a better indication of whole-plant change in growth (compared to length change) and does not assume exponential growth was occurring (growth rate constant makes this assumption). Two independent variables are investigated: 1) log(mean sulfide) and 2) log(mean initial sulfide).

General 4-Parameter Logistic Model

- $y = A + (B-A)/(1+e^{((C-x)/D)})$
- A = left-side horizontal asymptote (max response)
- B = right-side horizontal asymptote (min response)
- C = x value at the inflection point of the curve.
- D = slope parameter, indicating the speed with which the curve rises between min and max responses.

EC50 and EC20

- EC20* and EC50* are calculated for positive values of y. EC20* = 0.8A. EC50* = 0.5A.
- EC50* is a more statistically robust estimate than EC20*.
- Additional work is required to calculate the error around these estimates.

Model 1**Data set = Range Finder****Y = weight change****X = log(mean sulfide concentration)**

Formula: weight_change_mgd.1 ~ SSfpl(log_meansulfide_ugL, a, b, c, d)

Parameters:


Estimate	Std. Error	t value	Pr(> t)
a	0.49692	0.03914	12.696 6.50e-08 ***
b	-0.10906	0.08163	-1.336 0.2085
c	2.55198	0.15504	16.460 4.27e-09 ***
d	0.32791	0.11407	2.875 0.0151 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 '' 1

Residual standard error: 0.0712 on 11 degrees of freedom

Number of iterations to convergence: 0

Achieved convergence tolerance: 5.048e-06

Range Finder

Model 2**Data set = Definitive 1****Y = weight change****X = log(mean sulfide concentration)**

Formula: weight_change_mgd.1 ~ SSfpl(log_meansulfide_ugL, a, b, c, d)

Parameters:


Estimate	Std. Error	t value	Pr(> t)
a 0.43796	0.05032	8.704	2.90e-06 ***
b -0.26621	0.14325	-1.858	0.0901.
c 2.48534	0.18674	13.309	3.99e-08 ***
d 0.29275	0.11990	2.442	0.0327 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 '' 1

Residual standard error: 0.08906 on 11 degrees of freedom

Number of iterations to convergence: 0

Achieved convergence tolerance: 4.934e-06

Definitive 1

Model 3**Data set = Definitive 2****Y = weight change****X = log(mean sulfide concentration)**

Formula: weight_change_mgd.1 ~ SSfpl(log_meansulfide_ugL, a, b, c, d)

Parameters:


Estimate	Std. Error	t value	Pr(> t)
a 0.26964	0.02459	10.966	2.92e-07 ***
b -0.03783	0.02685	-1.409	0.1865
c 2.32803	0.76367	3.048	0.0111 *
d 0.05737	0.35484	0.162	0.8745

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 '' 1

Residual standard error: 0.05889 on 11 degrees of freedom

Number of iterations to convergence: 0

Achieved convergence tolerance: 5.639e-06

Definitive 2

Model 4**Data set = Range Finder****Y = weight change****X = log(mean initial sulfide concentration)**

Formula: weight_change_mgd.1 ~ SSfpl(log_meaninitialsulfide_ugL, a, b,
c, d)

Parameters:


Estimate	Std. Error	t value	Pr(> t)
a 0.48265	0.03612	13.364	3.82e-08 ***
b -0.08572	0.05944	-1.442	0.1771
c 2.72890	0.09719	28.079	1.37e-11 ***
d 0.22054	0.07586	2.907	0.0143 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 '' 1

Residual standard error: 0.07411 on 11 degrees of freedom

Number of iterations to convergence: 8

Achieved convergence tolerance: 7.537e-06

Range Finder

Model 5**Data set = Definitive 1****Y = weight change****X = log(mean initial sulfide concentration)**

Formula: weight_change_mgd.1 ~ SSfpl(log_meaninitialsulfide_ugL, a, b,
c, d)

Parameters:


Estimate	Std. Error	t value	Pr(> t)
a 0.41524	0.04717	8.804	2.60e-06 ***
b -0.21105	0.08016	-2.633	0.0233 *
c 2.62819	0.08731	30.102	6.43e-12 ***
d 0.16320	0.06513	2.506	0.0292 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 '' 1

Residual standard error: 0.09422 on 11 degrees of freedom

Number of iterations to convergence: 0

Achieved convergence tolerance: 5.564e-06

Definitive 1

Model 6**Data set = Definitive 2****Y = weight change****X = log(mean initial sulfide concentration)**

Formula: weight_change_mgd.1 ~ SSfpl(log_meaninitialsulfide_ugL, a, b,
c, d)

Parameters:

Estimate	Std. Error	t value	Pr(> t)
a 0.26937	0.02407	11.192	2.37e-07 ***
b -0.03867	0.03292	-1.175	0.265
c 2.57251	0.26466	9.720	9.81e-07 ***
d 0.04166	0.11854	0.351	0.732

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 '' 1

Residual standard error: 0.05907 on 11 degrees of freedom

Number of iterations to convergence: 21

Achieved convergence tolerance: 9.047e-06

Definitive 2